

Development, implementation and assessment of a concussion education programme for high school student-athletes

Jeffrey G. Caron^a, Scott Rathwell^b, J. Scott Delaney^{c,d}, Karen M. Johnston^{e,f}, Alain Ptito^{g,h,i} and Gordon A. Bloom^a

^aDepartment of Kinesiology and Physical Education, McGill University, Montreal, QC, Canada; ^bSchool of Human Kinetics, University of Ottawa, Ottawa, ON, Canada; ^cDepartment of Emergency Medicine, McGill University Health Centre, Montreal, QC, Canada; ^dMcGill Sports Medicine Clinic, Montreal, QC, Canada; ^eDivision of Neurosurgery, University of Toronto, Toronto, ON, Canada; ^fAthletic Edge Sports Medicine, Toronto, ON, Canada; ^gDepartment of Psychology, McGill University Health Centre, Montreal, QC, Canada; ^hMcGill University Health Centre Research Institute, Montreal, QC, Canada; ⁱMontreal Neurological Institute, Montreal, QC, Canada

ABSTRACT

Although experts have noted that adolescent athletes should be educated about concussions to improve their safety, there is no agreement on the most effective strategy to disseminate concussion education. The purpose of this study was to develop, implement and assess a concussion education programme. More precisely, four interactive oral presentations were delivered to high school student-athletes ($N = 35$, $M_{age} = 15.94$, $SD = 0.34$) in a large urban centre. Participants completed a questionnaire at three time-points during the season to measure changes in their knowledge (CK) and attitudes (CA) of concussions, and focus group interviews were conducted following the concussion education programme. Questionnaire data revealed participants' post-intervention CK scores were higher than their pre-intervention scores. During the focus groups, the student-athletes said they acquired CK about the role of protective equipment and symptom variability, and in terms of CA, they intended to avoid dangerous in-game collisions in the future. Our study was the first to create and deliver a concussion education intervention across multiple time-points, and to use mixed-methods in its assessment. These findings may be of interest to researchers, practitioners and stakeholders in sport who are invested in making the sport environment safer through concussion education and awareness.

ARTICLE HISTORY

Accepted 18 December 2016

KEYWORDS

Concussion education;
intervention; adolescent;
mixed method; knowledge
translation; injury
prevention

Approximately 54% of Canadians aged 15–19 participate in organised sport each year (Canadian Heritage, 2013). Sport participation has long been advocated as a way for youth and adolescents to acquire important life skills such as leadership and teamwork and to contribute to their physical and psychosocial well-being (Côté, Bruner, Erickson, Strachan, & Fraser-Thomas, 2010). However, adolescents who participate in high school athletics commonly suffer musculoskeletal injuries (e.g., Swensen et al., 2013). Concussions are one type of injury that has an elevated incidence rate in high school athletics (Marshall, Guskiewicz, Shankar, McCrea, & Cantu, 2015). In fact, researchers have found that concussions are particularly problematic for high school athletes because they tend to underreport symptoms (Register-Mihalik et al., 2013) and because their symptoms are more severe and persistent compared to older athletes (Williams, Puetz, Giza, & Broglio, 2015). Given that researchers are now linking multiple

statistics likely underestimate the true occurrence of concussions because signs of the injury (i.e., transient loss of consciousness) are rarely observable after acute injury (McCroly et al., 2013). As a result, many athletes are not evaluated for a concussion because they do not seek medical care for reasons that range from deliberately hiding symptoms from teammates, coaches and/or medical professionals, to inadvertently ignoring concussion symptoms due to a lack of knowledge about the injury (e.g., Davies & Bird, 2015; Delaney, Lamfook, Bloom, Al-Kashmiri, & Correa, 2015; Kroshus, Baugh, Daneshvar, & Viswanath, 2014; Register-Mihalik et al., 2013). For example, Delaney et al. (2015) found that student-athletes did not report concussion symptoms because they did not believe they had suffered a serious enough injury to warrant medical attention. Underreporting concussions has been a consistent finding across studies, despite the fact that large-scale concussion education initiatives such as the US